

On the Classification of quantum group structures on the group $GL(2)$

A. Aghamohammadi ^{1,2,*}, M. Khorrami ^{1,3,4}, A. Shariati ^{1,4}

¹ *Institute for Studies in Theoretical Physics and Mathematics, P. O. Box 5746, Tehran 19395, Iran.*

² *Department of Physics, Alzahra University, Tehran 19834, Iran.*

³ *Department of Physics, Tehran University, North-Kargar Ave. Tehran, Iran.*

⁴ *Institute for Advanced Studies in Basic Sciences, P. O. Box 159, Gava Zang, Zanzan 45195, Iran.*

* *E-Mail: mohamadi@irearn.bitnet*

Abstract

All quantum group structures on the group $GL(2)$ are classified. It is shown that there are only two such structures, the well known quantum groups $GL_{qp}(2)$ and $GL_{hh'}(2)$.

Although the group $GL(2)$ is not a complicated one, and its dimension is not so high, it seems that there are still some ambiguities in the classification of the quantum group structures on $GL(2)$.

There are two well-known deformations of this group [1]: the two parametric Drinfeld-Jimbo deformation $GL_{q,p}(2)$ [2], and the two parametric Jordanian deformation $GL_{h,h'}(2)$ [3]. It has been shown [4] that these two parametric deformations are related to each other by a singular transformation (contraction). It has been proved that, up to isomorphisms, there are only two quantum group structures on $GL(2)$ with a central quantum determinant, $GL_q(2)$ and $GL_h(2)$ [5]. Recently Kupershmidt has tried to classify all quantum group structures on the group $GL(2)$ [6]. He has concluded that there are three different quantum group structures on $GL(2)$: $GL_{q,p}(2)$, $GL_{h,h'}(2)$, and the new structure $GL_{h,q}(2)$. Here we will show that this new structure is, in fact, not new: by a suitable regular transformation, it falls in one of the two well-known structures $GL_{q,p}(2)$ or $GL_{h,h'}(2)$, depending on the deformation parameters. So, in fact there are only two non-equivalent quantum group structures on $GL(2)$.

To begin, we introduce the quantum-plane relations of $GL_{q,p}(2)$ and $GL_{h,h'}(2)$. For $GL_{q,p}(2)$ we have

$$\begin{aligned} xy &= qyx, \\ \xi^2 &= 0, \\ \eta^2 &= 0, \\ \eta\xi &= -p\xi\eta. \end{aligned} \tag{1}$$

And for $GL_{h,h'}(2)$

$$\begin{aligned} [x, y] &= hy^2, \\ \xi^2 &= h'\xi\eta, \\ \eta^2 &= 0, \\ \eta\xi &= -\xi\eta. \end{aligned} \tag{2}$$

Here x and y are the coordinates of the quantum plane, $\xi = dx$, and $\eta = dy$. The relations of the new quantum plane introduced in ref. [6] are

$$\begin{aligned} xy &= yx, \\ \xi^2 &= h'_0\xi\eta, \\ \eta^2 &= r_0\xi\eta, \\ \eta\xi &= -p_0\xi\eta. \end{aligned} \tag{3}$$

In fact, this is not the parametrization used in ref. [6]. We have used this parametrization for later convenience.

As it has been noted in ref. [6], one can use a transformation to make $r = 0$. (Applying the transformation changes (h'_0, r_0, p_0) to (h', r, p) .) The point is that, by a similar transformation, one can make both h'

and r equal to zero. So that this new structure is, in fact, the old $\text{GL}_{q,p}(2)$ with $q = 0$. There is, however, one exception: if $(p_0 - 1)^2 = 4h'_0 r_0$, one cannot make both h'_0 and r_0 equal to zero. Instead, one can make r equal to zero, and p equal to one. But this is $\text{GL}_{h,h'}(2)$ with $h = 0$. Here we present the above-mentioned transformations.

I Case $(p_0 - 1)^2 \neq 4h'_0 r_0$.

Using the linear transformation

$$\begin{pmatrix} \bar{\xi} \\ \bar{\eta} \end{pmatrix} = \begin{pmatrix} 1 & t_1 \\ 1 & t_2 \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad (4)$$

where t_i 's are the roots of the equation

$$r_0 t^2 + (1 - p_0)t + h'_0 = 0, \quad (5)$$

one can see that

$$\bar{\xi}^2 = \bar{\eta}^2 = 0, \quad (6)$$

and

$$\begin{aligned} \bar{\xi}\bar{\eta} &= (h'_0 + t_2 - p_0 t_1 + r_0 t_1 t_2)\xi\eta, \\ \bar{\eta}\bar{\xi} &= (h'_0 + t_1 - p_0 t_2 + r_0 t_1 t_2)\xi\eta. \end{aligned} \quad (7)$$

Note that, $(p_0 - 1)^2 \neq 4h'_0 r_0$ guarantees that $t_1 \neq t_2$, and this shows that the transformation (4) is nonsingular. Also note that the coefficients of $\xi\eta$ in (7) cannot be both zero; otherwise, from (6) and (7), one would conclude that $\xi^2 = \eta^2 = \xi\eta = \eta\xi = 0$. So, in this case, the structure is equivalent to $\text{GL}_{q=1,p}(2)$, where

$$p = -\frac{h'_0 + t_1 - p_0 t_2 + r_0 t_1 t_2}{h'_0 + t_2 - p_0 t_1 + r_0 t_1 t_2}. \quad (8)$$

II Case $(p_0 - 1)^2 = 4h'_0 r_0$.

In this case, the transformation (4) is singular. So we use

$$\begin{pmatrix} \bar{\xi} \\ \bar{\eta} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & t \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad (9)$$

where t satisfies (5), that is

$$t = \frac{p_0 - 1}{2r_0} = \frac{2h'_0}{p_0 - 1}. \quad (10)$$

Now we have

$$\begin{aligned} \bar{\eta}^2 &= 0, \\ \bar{\xi}^2 &= \frac{h'_0}{h'_0 + t}\bar{\xi}\bar{\eta}, \\ \bar{\eta}\bar{\xi} &= -\bar{\xi}\bar{\eta}. \end{aligned} \quad (11)$$

We see that, in this case, the structure is equivalent to $\text{GL}_{h=0,h'}(2)$ with

$$h' = \frac{h'_0}{h'_0 + t}. \quad (12)$$

These linear transformations on the quantum plane induce similarity transformations on the quantum matrix. So we have concluded that there are exactly two distinct deformations of $\text{GL}(2)$.

References

- [1] H. Ewen, O. Ogievetsky, & J. Wess; Lett. Math. Phys. **22** (1991) 297.
- [2] Yu. I. Manin; *Topics in noncommutative geometry*; Princeton, NJ: Princeton University Press.
- [3] A. Aghamohammadi; Mod. Phys. Lett. **A8** (1993) 2607.
- [4] A. Aghamohammadi, M. Khorrami, & A. Shariati; J. Phys. **A28** (1995) L225.
- [5] B. A. Kupershmidt; J. Phys. **A25** (1992) L1239.
- [6] B. A. Kupershmidt; J. Phys. **A27** (1994) L47.